Top DNA Polymerase

[Cat. No.] E - 3100-2 (500 units) E - 3101-2 (2000 units)

[Lot. No.] 0701

[Concentration] 5 units/uL

• Description : Bioneer's Top DNA Polymerase is isolated from recombinant E.coli strain containing the DNA polymerase gene from Thermus thermophilus. It exhibits highest activity at pH 9.0 and 72 °C.

• Applications : Polymerase Chain Reaction(PCR), Primer extension, TA cloning, gene sequencing.

Supplied with Enzyme

- 10X Reaction Buffer (1 mL) : 10X reaction buffer and a separate tube of 20 mM MgCl₂

- Dilution Buffer for Top DNA Polymerase (1 mL): 20 mM Tris-HCl, 0.5 mM EDTA,

1 mM DTT, 100 mM KCl, Stabilizers, 50 % Glycerol, pH 8.0

• Storage Condition : 20 mM Tris-HCI, 0.5 mM EDTA, 1 mM DTT, 100 mM KCI, Stabilizers, 50 % Glycerol pH 8.0, store at -20°C

• Unit Definition : One unit is defined at the amount of enzyme that will incorporates 10 nmol of dNTP into acid-insoluble material in 30 minutes at 72 °C.

• Quality Assurance : Nuclease activity is not detected after incubation of 1 ug of substrate DNA supercoiled plasmid and lambda/Hind III DNA - with 5 units of Top DNA Polymerase in 50 uL reaction volume with the supplied Reaction buffer for 18 hr at 37 °C and 70 °C.

- General Reaction Condition [20 uL reaction volume]
- Reaction mixture

Template* Primer (forward) Primer (reverse) 10X reaction buffer 10MM dNTPs mix. <i>Top</i> DNA Polymerase D.W	variable 5 ~ 10 pmoles 5 ~ 10 pmoles 2 uL variable (0.3 ~ 1.0 mM) 0.5 ~ 2.0 unit variable	* Amounts of template Plasmid and lambda DNA \rightarrow more than 1 pg Bacterial genomic DNA \rightarrow more than 10 pg Human genomic DNA \rightarrow more than 1ng
Total	20 uL	

Total

Notice to purchaser

This enzyme is specifically optimized for increasing base incorporation rate by inactivating 5'->3' exonuclease activity. Therefore, this is not recommended to use for Real Time PCR using Tagman[®] probe, another product for which will be released soon.

Ordering Information

[Related BIONEER 's Top DNA Polymerase Products]

Cat. No.	Products	
E-3100 – 1	Top DNA Polymerase 500 units, 10 mM dNTPs, 10X reaction buffer and a separate tube of 20 mM $MgCl_2$	
E-3101 – 1	Top DNA Polymerase 2,000 units, 10 mM dNTPs, 10X reaction buffer and a separate tube of 20 mM $MgCl_2$	
E-3100 – 3	Top DNA Polymerase 500 units, 10X reaction buffer and a separate tube of 20 mM $MgCl_2$	
E-3101 – 3	Top DNA Polymerase 2,000 units, 10X reaction buffer and a separate tube of 20 mM MgCl ₂	
E-3100	Top DNA Polymerase 500 units, 10 mM dNTPs , 10X reaction buffer with MgCl_2	
E-3101	Top DNA Polymerase 2,000 units, 10 mM dNTPs , 10X reaction buffer with $MgCl_2$	
D-3001	10 mM dNTP mixture (1.0 mL), each dNTP 2.5mM	
EB-1002	10X reaction buffer with MgCl ₂	

Note

For research use only. Not for use in diagnostic or therapeutic procedures.

Bioneer shall not in any event be liable for incidental or special damage of any kind resulting from any use except on application(s).

If you use short length primer or random primer in PCR, you may detect unexpected PCR product (or non-specific bands)